

Durham Research Online

Deposited in DRO:

26 March 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Harper, D.A.T. (2019) 'The Golden Spike still glitters : the (re)construction of a global chronostratigraphy.', *Acta Geologica Sinica : Supplement.*, 93 (S3). pp. 24-27.

Further information on publisher's website:

<https://doi.org/10.1111/1755-6724.14234>

Publisher's copyright statement:

This is the accepted version of the following article: Harper, D.A.T. (2019). The Golden Spike Still Glitters: The (Re)construction of a Global Chronostratigraphy. *Acta Geologica Sinica* 93 (supplement)(S3): 24-27, which has been published in final form at <https://doi.org/10.1111/1755-6724.14234>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Additional information:

Published on behalf of the Geological Society of China.

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

The Golden Spike Still Glitters: The (Re)construction of a Global Chronostratigraphy

David A.T. HARPER

Palaeoecosystems Group, Department of Earth Sciences, Durham University, Durham DH1 3LE, UK

Corresponding author's E-mail: david.harper@durham.ac.uk

Introduction

Stratigraphy, placing the sequence of events in Earth history into order, is a fundamental part of the geosciences. Traditionally the starting point is a regional stratigraphy, based on lithological, mappable units, correlated by fossils with local, regional and global chronostratigraphical schemes. Chronostratigraphic units were traditionally defined on their fossil content, as essentially unit stratotypes. But, commonly there were no clear boundaries, the lower and upper parts of units being coincident with major faunal changes such as extinction events that delimited given units. There were problems with this approach and during the latter part of the 1900s, geologists moved towards a more rigorous and robust method of defining chronostratigraphic units based on Global Stratotype Sections and Points (GSSPs), anchored by 'Golden Spikes'. As the stratigraphic column reaches completion with the majority of the stages, series and systems defined by GSSPs, the challenge is now to calibrate the geological timescale and refine the correlations of regional and global stratigraphical schemes. Radiometric dating, isotope curves, magnetostratigraphy and orbital-tuning are providing more accurate and precise methods for correlation, and dating the timescale in absolute terms.

The birth of stratigraphy

Pioneer work by the Danish polymath Nicolaus Steno (1638-1686) in northern Italy, during the late 17th Century, established the obvious fact that older rocks are overlain by younger rocks, if the sequence has not been inverted. His law of superposition of strata is fundamental to all stratigraphic studies. In addition, Steno established in experiments that sediments are deposited horizontally and rock units can be traced laterally, often for considerable distances; remarkably simple concepts to us now, but paradigm shifting at the time. This, however, was preceded by the studies of Leonardo da Vinci (1452–1519) who essentially rediscovered geological perspective, some 200 years before Steno, during the Renaissance. In his drawing of the hills of Tuscany, da Vinci portrayed a clear sequence of laterally continuous, horizontal strata displaying the concept of superposition; he also noted fossil shell beds in some of his sketches.

The organization of rocks

About a century after Steno, Giovanni Arduino recognized, again using superposition, three basically different rock suites in the Italian part of the Alpine belt. A crystalline basement of older rocks, deformed during the Late Palaeozoic Variscan orogeny, was overlain unconformably by mainly Mesozoic limestones deformed later during the Alpine orogeny; these in turn were overlain unconformably by poorly consolidated clastic rocks, mainly conglomerates. These three units constituted his primary, secondary and tertiary systems; the last term is still used by some for the interval of geological time succeeding the Cretaceous. These three divisions were used widely to describe rock successions elsewhere in Europe showing the same patterns, but these three systems were not necessarily the time correlatives of the type succession in the Apennines. But they needed a time frame. The concept of deep time was provided by James Hutton (1726-1797), while nearly a century later, the role of fossils in stratigraphy was established by William Smith in Britain, and Georges Cuvier and Alexandre Brongniart in France together with Albert Oppel in Germany. William Smith (1769–1839), in the course of his work as a canal engineer in England, realized that different rock units were characterized by distinctive groups or assemblages of fossils. In a traverse from Wales to London, Smith encountered successively younger rocks, and he documented the change from the trilobite-dominated assemblages of the Lower Palaeozoic of Wales through Upper Palaeozoic sequences with corals and thick Mesozoic successions with ammonites; finally, he reached the molluscan faunas of the Tertiary strata of the London Basin. In France, a little later, the noted anatomist Georges Cuvier (1769-1832) together with Alexandre Brongniart (1770–1849), a leading mollusc expert of the time, ordered and correlated Tertiary strata in the Paris Basin using series of mainly terrestrial vertebrate faunas, occurring in sequences separated by supposed biological catastrophes. Distinctive lithologies and biotic assemblages would now allow the recognition of a sequence of major stratigraphical units, the systems.

The scramble for systems

Geological time was divided up by the efforts of pioneering British, French and German geologists between 1790 and 1840. The divisions were made first for practical reasons – one of the first systems to be named was the Carboniferous ("coal-bearing"), a unit of rock that early industrialists were keen to exploit. The Cretaceous was established based on the widespread

distribution of chalk whereas the Triassic in northern Europe could be split into three distinctive megafacies. Many of the others were based on geographic areas where the systems cropped out or in the case of the Lower Palaeozoic, the proximity to the home of some ancient Welsh tribes.

Some systems were established not without some controversy and rancor. During the 1830s, Roderick Murchison (1792–1871) and Adam Sedgwick (1785–1873) collaborated, and then clashed, over the Lower Palaeozoic. Sedgwick named the Cambrian and Murchison named the Silurian, based on sections in Wales. Each claimed the middle ground for his system, so what Murchison called the “Lower Silurian”, Sedgwick called “Upper Cambrian”. This territorial claim was resolved later by Charles Lapworth (1842–1920), who agreed with neither of them, and named the contentious rock successions the Ordovician in 1879. In fact, the Ordovician is one of the longest and most lithologically diverse of the geological systems but it was only formally accepted by the international community in 1960. By the mid-1800s a chronostratigraphic framework was in place permitting Darwin (1859) to use palaeontological data to illustrate his theory of evolution and John Phillips to explore the history of life on Earth (Phillips, 1860).

A new age in stratigraphy: The concept and implementation of GSSPs

It became clear, as greater precision was required to correlate events and strata globally, that many of the original boundaries of the geological systems were separated from each other by unconformities (Remane, 2003). For the pioneers, unconformities provided a convenient break between systems and, more importantly, it satisfied their view that the major divisions of Earth’s history should be divided by global, catastrophic events. Unfortunately, many of these unconformities turned out to be only regional breaks that occurred in Europe, but not elsewhere. The bases of many systems were represented by stratigraphic gaps, obviously, providing a poor basis for the global correlation of systemic boundaries.

A new approach was required and this was provided by the International Commission on Stratigraphy (ICS). All the system boundaries have been reinvestigated by working groups of the ICS, as have many of the series and stage divisions (Cohen et al., 2013). The potential of each base for international correlation must be maximized, and the bases of these systems must be placed within intervals of continuous sedimentation, with diverse and abundant faunas and floras in geographically and politically accessible areas that can be conserved and protected; ideally the sections should have escaped metamorphism and tectonism and have the potential to use non-biological proxies to aid correlation. The base of a chronostratigraphic interval is defined in a unique stratotype section, in a type area using the concept of a “golden spike” or marker point (Hedberg, 1976). All the usual criteria for a workable stratotype section must, of course, be satisfied (Cowie et al., 1986). The golden spike, which represents a point in the rock section and an instant in geological time, is then driven into the section, at least in theory (Holland, 1986). In reality, the spike is usually adjusted to coincide with the first appearance (FAD) of a distinctive, recognizable fossil within a well-documented lineage, though non-biological markers are also important. The ranges of all fossils occurring across the boundary, together with geochemical and other proxies, are documented in detail as aids to correlating within the section and with sections elsewhere. Establishing stratotypes and golden spikes requires international agreement through the various working groups and subcommissions of the ICS, and that can sometimes be time-consuming and hard to achieve. But, when finally ratified by the IUGS, this horizon will then be the Global Stratotype Section and Point (GSSP) for this unit and reported on the regularly-updated ICS timescale chart (Cohen et al., 2018; Figure 1).

Calibrating the geological timescale

Exciting advances in technology are delivering an improved calibration and correlation of strata; both regional and global-scale divisions at high resolutions can be better linked and the rates of biological and geological processes assessed with greater accuracy. Biological data are commonly matched by chemostratigraphy, cyclostratigraphy and magnetostratigraphy together with high precision absolute dates produced by techniques using new generations of mass spectrometers, sometimes with an accuracy of some 10 kyr. Successions can also be orbitally-tuned using Milankovitch cycles based on eccentricity (variation in the shape of the Earth’s orbit from nearly circular to elliptical; 100 kyr cycle), obliquity (wobble of the Earth’s axis; 41 kyr cycle) and precession (change in direction of the Earth’s axis relative to the sun; 23 kyr cycle). Throughout the stratigraphic record, there are many successions of rhythmically alternating sedimentary beds, for example repeated pairs of limestone and calcareous shales that may have been controlled by Milankovitch processes, although not all modulations may have been constant in deep time. These data, biological and non-biological, can be integrated in the definition and description of chronostratigraphic units. The recent establishment of the GSSP for the Chattian Stage (Paleogene System) is an elegant exemplar (Coccioni et al., 2018).

It is now possible to provide such integrated stratigraphies for an entire country. Shen and Rong (2019) have amassed and integrated all available stratigraphic data from China and compiled these from the Ediacaran to Quaternary in a landmark publication. The reconstruction of the geological timescale is only the start.

Key words: Stratigraphy, biostratigraphy, GSSPs, Milankovitch cycles, timescale

Acknowledgements: Harper thanks the Leverhulme Trust for financial support. This is a contribution to IGCP 653, ‘The onset of the Great Ordovician Biodiversification Event’.

References

- Coccioni, R., Montanari, A., Bice, D., Brinkhuis, H., Deino, A., Frontalini, F., Lirer, F., Maiorano, P., Monechi, S., Pross, J., Rochette, P., Sagnotti, L., Sideri, M., Sprovieri, M., Tateo, F., Touchard, Y., Van Simaey, S., and Williams, G.L., 2018. The Global Stratotype Section and Point (GSSP) for the base of the Chattian Stage (Paleogene System, Oligocene Series) at Monte Cagnero, Italy. *Episodes* 41, 17–32.
- Cohen, K.M., Finney, S.C., Gibbard, P.L., and Fan Junxuan, 2013. The ICS International Chronostratigraphic Chart. *Episodes* 36, 199–204.
- Cohen, K.M., Harper, D.A.T., and Gibbard, P.L., 2018. *International Chronostratigraphic Chart*. International Commission of Stratigraphy (www.stratigraphy.org).
- Cowie, J.W., Ziegler, W., Boucot, A.J., Bassett, M.G., and Remane, J., 1986. Guidelines and statutes of the International Commission on Stratigraphy (ICS). *Courier Forschungsinstitut Senckenberg* 83, 1–14.
- Darwin, C., 1859. *On the origin of species by means of natural selection, or the preservation of favoured races in the struggle for life*. John Murray, London.
- Hedberg, H. D. (ed.) 1976. *International Stratigraphic Guide: a Guide to Stratigraphic Classification, Terminology, and Procedure*. International Subcommission on Stratigraphic Classification of IUGS Commission on Stratigraphy. John Wiley and Sons, New York.
- Holland, C.H. 1986. Does the Golden Spike still glitter? *Journal of the Geological Society, London* 143, 3–21.
- Phillips, J., 1860. *Life on Earth: its origin and succession*. University Press, Cambridge.
- Remane, J. 2003. Chronostratigraphic correlations: their importance for the definition of geochronologic units. *Palaeogeography, Palaeoclimatology, Palaeoecology* 196, 7–18.
- Shen Shuzhong and Rong Jiayu (eds) 2019. Integrative stratigraphy and timescale of China. *Science China: Earth Sciences* 62, 1–348.

About the first author (also the corresponding author)

HARPER, David A.T., Male, born in 1953 in Edinburgh; PhD, DSc; graduated from Imperial College, London and Queen's University, Belfast; Professor of Palaeontology in Earth Sciences and Principal of Van Mildert College, Durham University, UK. He is now researching some of the major events in the history of life, for example the Cambrian Explosion, Great Ordovician Biodiversification Event and the end Ordovician mass extinctions. His research programmes have taken him to many parts of the world including Australia, Chile, China, Greenland, Russia, Svalbard and Tibet.

Email: david.harper@durham.ac.uk; phone: +44 01913347143.

Fig. 1. The International Chronostratigraphic Chart, 2018, version 8 (Cohen et al., 2018).



International Commission on Stratigraphy

v 2018/08



Units of all ranks are in the process of being defined by Global Boundary Stratotype Section and Points (GSSP) for their lower boundaries, including those of the Archean and Proterozoic, long defined by Global Standard Stratigraphic Ages (GSSA). Charts and detailed information on ratified GSSPs are available at the website <http://www.stratigraphy.org>. The URL to this chart is found below.

Numerical ages are subject to revision and do not define units in the Phanerozoic and the Ediacaran; only GSSPs do. For boundaries in the Phanerozoic without ratified GSSPs or without constrained numerical ages, an approximate numerical age (~) is provided.

Ratified Subseries/Subepochs are abbreviated as U/L (Upper/Late), M (Middle) and L/E (Lower/Early). Numerical ages for all systems except Quaternary, upper Paleogene, Cretaceous, Triassic, Permian and Precambrian are taken from 'A Geologic Time Scale 2012' by Gradstein et al. (2012), those for the Quaternary, upper Paleogene, Cretaceous, Triassic, Permian and Precambrian were provided by the relevant ICS subcommissions.

Colouring follows the Commission for the Geological Map of the World (<http://www.ccgm.org>)

Chart drafted by K.M. Cohen, D.A.T. Harper, P.L. Gibbard, J.-X. Fan
(c) International Commission on Stratigraphy, August 2018

To cite: Cohen, K.M., Finney, S.C., Gibbard, P.L. & Fan, J.-X. (2013; updated)
The ICS International Chronostratigraphic Chart. Episodes 36: 199-204.

URL: <http://www.stratigraphy.org/ICSchart/ChronostratChart2018-08.pdf>